



Graduated Medical Compression Stockings

Date: _____

Patient Name: _____

Diagnosis: _____

Physician Signature: _____ **(Dispense as written no substitution)**

Practice #: _____ Dispense Quantity: _____ Rep: _____

Please indicate Compression and Style for Compression garment.

RX COMPRESSION (select one)

- | | | |
|--|---|--|
| <input type="checkbox"/> 15-20 mmHg | <input type="checkbox"/> Class I: 20-30 mmHg | <input type="checkbox"/> Class II: 30-40 mmHg |
| <input type="checkbox"/> Anti-embolism (Non ambulatory) | <input type="checkbox"/> Class III: 40-50 mmHg | <input type="checkbox"/> Class IV: 50-60 mmHg |

NON ELASTIC COMPRESSION

- CircAid Juxtalite® Venous**
 CircAid Juxtafit® Lymphedema

PNEUMATIC COMPRESSION

- Venous**
 Lymphedema

(select one style form chart)

STYLE



	Calf	Thigh	Panty	Compression sleeve		Glove	
	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> Left	<input type="checkbox"/> Right	<input type="checkbox"/> Left	<input type="checkbox"/> Right

Dealer: _____

PHYSICIAN'S THERAPY GUIDE

INDICATIONS

SUPPORT

- Slight varicosities
- Tired aching legs

RX COMPRESSION

CLASS I: 20-30 mmHg

- Heaviness & fatigue in legs
- Minor varicosities
- Initial varicosities during pregnancy
- Post surgical
- Minor ankle, leg & foot swelling
- Post sclerotherapy
- Superficial thrombophlebitis
- Prophylactic treatment during pregnancy
- Hereditary disposition toward vein disease

CLASS II: 30-40 mmHg

- Moderate varicosities
- Moderate edema
- Lymphatic edema
- Management of venous ulcers
- Prevention of ulcer recurrence
- Prevention of post- thrombotic syndrome
- Orthostatic hypotension
- Post surgical
- Post sclerotherapy
- Effective scar formation after burns
- Pregnancy with previous phlebitis
- Stasis dermatitis due to CVI

CLASS III: 40-50 mmHg

- Severe varicosities
- Severe edema
- Lymphatic edema
- Management of venous ulcers
- Post phlebotic syndrome
- Chronic venous insufficiency
- Resersible lymphedema
- Venous ulcers
- Post surgery
- Orthostatic hypotension

CONTRAINDICATIONS:

According to the guideline of the German Phlebological Society:

Absolute Contraindications: Advanced peripheral arterial occlusive disease, decompensated heart disease, septic phlebitis, phlegmasia coerulea dolens, congestive heart failure, cutaneous infections, severe arterial insufficiency. **Relative contraindications:** suppurative dermatoses, intolerance of compression stocking fabric, sensory disturbances of the limb, advanced peripheral neuropathy, primary chronic arthritis.

Medi assumes no liability as a result of any contraindicated use of this product.